


# *In-situ* Detection and Mapping of Microbial Life in Ice with a Miniaturized Biospectrologger

P. B. Price, N. Bramall, R. Bay (U. C. Berkeley)  
W. Hug (Photon Systems)

Despite growth curves in pure culture, and despite statements in the literature that the limit for life is  $\sim -20^{\circ}\text{C}$ , I will show that it is worthwhile to search for living microbes in cold planets.



# Unusual route to invention



AMANDA high-energy neutrino observatory ~1993



Dust in glacial ice governs its optical properties ~1994



Brainstorming for proposed NSF STC “Deep Ice” ~1998



Microbes in veins in ice ~2000



Dust logger ~2001




Biospectral logger (BSL)



Discoveries?

Jumps in dust signal in 3054 m borehole (Greenland)  $\Rightarrow$  abrupt climate changes. Do microbes accompany dust and volcanic ash?



## Technology Goal

- Construct biospectral logger (miniBSL) to fit into a 5-cm borehole; 224 nm laser; 6 notch filters + 1 ND filter; 7 PMTs; noninvasive

## Science Goals

- Log 300-m-deep borehole in glacial ice 8 km from S. Pole. On Jan. 16, N. Bramall logged this borehole with a large BSL.
- With Peter Doran *et al.*, log 19-m ice cover, microbial mats, and hypersaline water in Lake Vida, Antarctica.
- Scan ice cores at NICL ; do microbes correlate with climate changes, basal ice, volcanic ash, age and T of ice? Do they live in veins in ice?
- From fluorescence spectra, identify types, live/dead ratio, microhabitats, water/ice ratio in subsurface,...

## Microbial life on Earth “follows the water”




- Ice has microbes in interconnecting aqueous,  $\mu\text{m}$ -size veins (salt-rich on Mars; acid-rich on Europa)
- Permafrost has microbes on nm-thick films of unfrozen water at ice/grain surfaces.

# Melting points of some aqueous eutectics


HCl	-90 C
HNO <sub>3</sub>	-43 C
H <sub>2</sub> SO <sub>4</sub>	-73 C
Methanosulfonic acid	-75 C
Formaldehyde	-92 C
Formic acid	-49 C
NaCl · 2H <sub>2</sub> O	-22 C
CaCl <sub>2</sub> · 6H <sub>2</sub> O	-50 C
MgCl <sub>2</sub> · 12H <sub>2</sub> O	-33 C
CaCl <sub>2</sub> · 6 H <sub>2</sub> O + MgCl <sub>2</sub> · 12H <sub>2</sub> O	-55 C
NH <sub>4</sub> OH	-84 C
NH <sub>4</sub> CN	-100 C


Direct evidence that adapted microorganisms can exist and metabolize in a network of liquid veins at T down to  $-90^{\circ}\text{C}$ .

Fluorescing microbe in a vein in sea ice at  $-15^{\circ}\text{C}$  (K. Junge and J. Deming)










Water film thickness at grain/ice Interface depends on temperature and rock type: clay minerals are best: high surface to volume


Microbes live on mineral surfaces in permafrost and extract energy from the thin films that remain liquid even at low temperature.

- blue = DAPI-stained bacteria
- red = chlorophyll fluorescence


(Priscu et al.: ice-covered Lake Bonney, Antarctica)




# GISP2 (Greenland) 3054 m core; silty ice in bottom 13 m







near bottom  
of Greenland ice  
core  $\Rightarrow$  microbes are  
metabolizing in ice.



V. Miteva finds  $\sim 7 \times 10^7$   
cells/cm<sup>3</sup> in deepest ice.

C. Tung (my undergrad)  
finds, on average,  $\sim 20$  cells

# Our geochemical method for inferring metabolic rate of trapped, dormant microbes at low temperature:

$\mu$  = metabolic rate per cell in gC gC<sup>-1</sup> h<sup>-1</sup>


$$\mu = Y_j / n_j m_j t$$

$Y_j$  = metabolic yield of gas of type  $j$ ;




$n_j, m_j$  = conc. and mass of microbes producing  $j$ ;

$t$  = retention time of reactants


# Nitrifying bacteria In Vostok ice



# SEM pictures of bacteria from Vostok ice core at 3593 m







# Microbe metabolism

- Activation energies  $\sim 110 \text{ kJ/mole}$
- Rates scale as  $10^6:10^3:1$
- Metabolism down to  $-40^{\circ}\text{C}$ ; no minimum
- In Vostok ice at  $-40^{\circ}\text{C}$ , 1 turnover per  $10^8 \text{ yr}$ !
- In “survival” mode, metabolic rate for repair  $\approx$  spontaneous damage rate.



Nathan Bramall and Ryan Bay with BSL-1 at Lake Tahoe. Bramall logged a South Pole borehole with a bigger version on Jan. 16.



Log of Lake Tahoe;  
BSL-1 excited at  
370 nm

Fluorescence from  
Chl-a and NADH  
→

NADH fluorescence  
spectra of known  
microbes compared  
with Tahoe  
microbes →



MiniBSL is a *discovery* instrument: design sensitivity  $\approx 1$  cell/cm<sup>3</sup>.






At 224 nm, microbial fluorescence dominates over mineral fluorescence. Examples:

- Chl-a  $\Rightarrow$  photosynthesis
- Trp  $\Rightarrow$  all cells
- Tyr  $\Rightarrow$  spore coats
- F420  $\Rightarrow$  a coenzyme in methanogens and haloarchaea
- Guanosine/adenosine in acid veins  $\Rightarrow$  live/dead test
- PAHs

To detect other biomolecules, use other wavelengths



- Fulvic and humic acids,...
- Bacteriorhodopsin  $\Rightarrow$  in some haloarchaea
- NADH, flavins,...

With ND filter, scattering w/o fluorescence  $\Rightarrow$  dust, volcanic ash,...



Focuses on borehole wall miniBSL

Focuses inside the ice





# Future potential applications


Beacon Valley (Antarctica): ~8 Myr-old permafrost/ice

Borehole in deep mine (anaerobes on walls)

Subglacial Lake Vostok (exotic microbes?)

Mars subsurface?



Warm diapirs in Europa's ice?



MiniBSL could search in a Martian borehole for biomolecules in ice, permafrost, or rock and for dust and ash in clean ice.

Below the impact-gardened depth (3-14 m), organic molecules and even microbial life might avoid oxidants and surface irradiation.





Does microbial  
life exist in  
veins within  
warm diapirs in  
European ice?